

UTTG-17-92

July 1992

Trivial Spectrum of Free 1+1 Light-Cone Strings*

Eric Smith

Theory Group
Department of Physics
University of Texas
Austin, Texas 78712

ABSTRACT

The BRST cohomology of 1+1 strings in a free light-cone gauge contains only the two-dimensional tachyon, and excludes all excited states of both matter and ghosts, including the special states that arise in the continuum conformal gauge quantization and in the $c = 1$ matrix models. This exclusion takes place at a very basic level, and therefore may signal some serious problems or at least unresolved issues involved in this gauge choice.

*Research supported in part by the Robert A. Welch Foundation and NSF Grant PHY 9009850

Fundamental strings used to be defined by a requirement that they possess only transverse physical excitations [1]. That changed with the discovery that Polyakov's critical string theory in $1 + 1$ dimensions possesses excited states of timelike oscillators as nontrivial representatives of its BRST cohomology at special values of the center of mass momentum [2]. Since then these states have been extensively studied [3, 4], and while their algebra is now well described [5], their physical interpretation remains obscure. It would be useful to have a description of these states in an easily generalizable light-cone gauge, to understand both how they were missed by earlier quantizations and whether they have any relevance in higher dimensional or effective string theories with similar backgrounds.

An attempt to formulate such a description was begun in [6]. It centered on a gauge which, in $1 + 1$ dimensions, leaves a free theory of its unfixed degrees of freedom, but possesses propagating ghosts and a nontrivial BRST charge. This attempt appears to fail, because unlike BRST quantization in conformal gauge, which admits many states to the mass shell but excludes almost all the excited ones from a reduced cohomology, the free light-cone gauge seems to exclude all excited states directly from the mass shell. This drastic exclusion takes place at a very basic level, and could signal some serious inconsistency of this gauge or of the implementation of it given in [6]. Because the program may have value, but the details remain problematic, this note describes the calculation of the spectrum.

The following results are brought forward from previous work: The starting point is a Liouville-type action

$$S_C = \frac{-1}{4\pi\alpha'} \int d^2\sigma \sqrt{g} \left(g^{ab} \partial_a X^\mu \partial_b X^\nu \eta_{\mu\nu} + \alpha' n \cdot X R^{(2)} \right) \quad (1)$$

with world-sheet coordinates (τ, σ) and metric g_{ab} describing a Euclidean cylinder of spatial (σ) period $-2\sqrt{2}\pi^1$, and target space light-cone coordinates $X^\pm \equiv (X^0 \pm X^1)/\sqrt{2}$ with flat target space metric $\eta_{\mu\nu}$ and n a constant target space vector. α' is the inverse string tension. Fadeev-Popov [7] imposition of the gauge condition

$$\hat{g}^{--} \equiv \sqrt{g} g^{--} = 0; \quad \sqrt{g} = 1; \quad \hat{X}^+ \equiv X^+ + \frac{\alpha' n^+}{2} \log \sqrt{g} = \tau \quad (2)$$

leaves the free actions

$$\begin{aligned} S_{C+M} &= \frac{1}{\pi} \int d^2\sigma \hat{g}^{++} \left[\left(1 - \frac{\partial_+}{M} \right) \left(\frac{\sqrt{2}}{4\alpha'} \hat{X}^-, + \right) + \frac{1}{4\alpha'} \right] \\ S_{GH} &= \frac{1}{\pi} \int d^2\sigma (wu_{,+} + zv_{,+}) \end{aligned} \quad (3)$$

for matter and ghosts, respectively. Here $\hat{g}^{++} \equiv \sqrt{g} g^{++}$, $\hat{X}^- = X^- - \frac{\alpha' n^+}{2} \log \sqrt{g} \equiv X^- - \frac{1}{\sqrt{2}M} \log \sqrt{g}$, and metric components and derivatives are indexed with respect to world sheet coordinates $\sigma^\pm \equiv (\tau \pm \sigma)/\sqrt{2}$. The ghost (antighost) components $u, v, (w, z)$ come from a decomposition of the reparametrization ghosts c^+ and c^- :

$$c^- = u; \quad c^+ = -(1 + \frac{\partial_-}{M})u + e^{-\sqrt{2}M\tau}v \quad (4)$$

¹This range is chosen so that the corresponding complex coordinates take the usual form.

and $\hat{X}^{+-}$ and $\hat{g}^{--}$ antighosts, named respectively b_f and $\hat{b}_{--}$:

$$b_f = -Me^{\sqrt{2}M\tau}z; \quad \hat{b}_{--} = \frac{w}{2} - \frac{1}{2M}e^{\sqrt{2}M\tau}z_{,-}. \quad (5)$$

The Weyl ghost is trivial and has already been integrated out. The subscript c_{+M} on the matter action indicates that the factor $\hat{g}^{++}/4\alpha'$ is a term extracted from the measure and added to the canonical action to give the resulting full matter action the symmetries of the usual Liouville theory[8].

The equations of motion in the matter sector are solved by a field decomposition of the form²

$$\hat{g}^{++} = f + e^{-\sqrt{2}M\tau}g - 2; \quad \frac{\sqrt{2}}{4\alpha'}\hat{X}^- = h - e^{\sqrt{2}M\tau}j - \frac{\sqrt{2}}{4\alpha'}\tau, \quad (6)$$

with f, g, h, j as well as u, v, w, z functions only of σ^- on shell. Boundary conditions for the cylinder require periodicity of all of these functions when $\sigma \rightarrow \sigma - 2\sqrt{2}\pi$. At this point it is convenient to take $\sigma = -is$ so that σ^- becomes the complex variable w , in terms of which the functions f, g, h, j, u, v, w, z are analytic. Because the currents of interest are all tensors, it is convenient to map the cylinder to a plane with coordinates $z = e^w$. Then the measure for path integration in this theory is specified by imposing the free field propagators

$$\langle vz' \rangle = \langle uw' \rangle = \langle hf' \rangle = \langle jg' \rangle = \frac{1}{z - z'}. \quad (7)$$

²The additive constants which appear here are a convenience which simplified the expression of the conserved currents in [6].

The reparametrization symmetries for matter and ghosts have no simple form in terms of fields $\theta u, \theta v$ (drawn from the ghost decomposition (4) by adding a Grassmann parameter θ to the fields u, v), but the associated conserved currents take a simple form on shell:

$$\begin{aligned} J_{uC} &= \left(-f \partial h - \partial g j - 2g \partial j + \frac{2}{M} \partial^2 h \right); \\ J_{vC} &= (m f j + 2 \partial j); \\ J_{uGH} &= (-v \partial z + u \partial w + 2 \partial u w); \\ J_{vGH} &= (u \partial z). \end{aligned} \tag{8}$$

The J_u s generate an analytic Virasoro algebra, and the J_v s an analytic $U(1)$. There are no antianalytic currents associated with these symmetries. The BRST charge Q takes the standard form:

$$Q = \frac{1}{2\pi i} \oint dz \left[u J_{uC} + v J_{vC} + \frac{1}{2} (u J_{uGH} + v J_{vGH}) \right]. \tag{9}$$

The conformal dimensions of the analytic fields are computed from their operator product expansions with the currents J_u , and take values $(1, 2, 0, -1; -1, 1, 2, 0)$ respectively for $(f, g, h, j; u, v, w, z)$. The currents J_{uC}, J_{uGH} have dimension 2 and central charge $+28, -28$ respectively, and the currents J_{vC}, J_{vGH} have dimension zero and no anomaly terms. Of the OPEs of J_u with the free fields, the only anomalous product is $J_{uC} f'$, which contains the term $4/(z - z')^3 M$.

Decomposition of the analytic fields into oscillators is done in the usual way, with

$$\alpha_n \equiv \frac{1}{2\pi i} \oint dz z^{n+d-1} \alpha(z); \quad \alpha(z) = \sum_n \frac{\alpha_n}{z^{n+d}}, \tag{10}$$

where α represents the name of the field and d is its conformal dimension. The special names $L_n, K_n; \lambda_n, \kappa_n$ will be given to the expansion coefficients in $J_{uC}, J_{vC}; J_{uGH}, J_{vGH}$, respectively.

The first point to notice about this theory is that the matter fields, and not just their derivatives, decompose directly into analytic fields and constants. The consequences of this decomposition become apparent when the generators of the current algebra are written in terms of oscillators, as:

$$\begin{aligned}
L_m &= \sum_n [(m-n) : f_n h_{m-n} : + (2m-n) : g_n j_{m-n} :] \\
&\quad + \frac{2}{M} m(m+1) h_m + \delta_m \\
\lambda_m &= \sum_n [(m-n) : v_n z_{m-n} : + (-m-n) : u_n w_{m-n} :] - \delta_m \\
K_m &= \sum_n [M f_n j_{m-n}] - 2(m-1) j_m \\
\kappa_m &= \sum_n [n z_n u_{m-n}]
\end{aligned} \tag{11}$$

and

$$\begin{aligned}
Q &= \sum_m [u_m L_{-m} + v_m K_{-m}] \\
&\quad + \sum_{m,n} \left[\frac{(m-n)}{2} : u_m u_n w_{-m-n} : + n u_{-m-n} : v_m z_n : \right] - u_0,
\end{aligned} \tag{12}$$

where $: :$ denotes creation-annihilation normal ordering, and the ordering constants in L_0, λ_0 and Q are fixed by the action of the commutators on the ground state, as implied by the current-current OPEs [9].

A standard technique for computing BRST cohomology is to write Q as

a sum

$$Q = u_0 (L_0 + \lambda_0) + v_0 (K_0 + \kappa_0) - 2w_0 \sum_{n>0} n u_{-n} u_n + \hat{Q} \quad (13)$$

and show how the cohomology of Q relates to that of $\hat{Q}$ and to the mass shell conditions that arise as a consequence of reparametrization invariance [9].

The ghost zero mode sectors of this theory are slightly more complicated than those for a simple Virasoro algebra, but it is still straightforward to show that every BRST cohomology class can be represented by a state $|\text{phys}\rangle$ for which

$$(L_0 + \lambda_0) |\text{phys}\rangle = 0; \quad (K_0 + \kappa_0) |\text{phys}\rangle = 0. \quad (14)$$

The elementary commutation relations implied by (7) make it possible to define number operators for the excitations which annihilate the ground state and satisfy

$$[N_{\alpha_n}, \alpha_{-m}^k] = k \alpha_{-m}^k \delta_{n-m}; \quad m > 0, \quad (15)$$

in terms of which

$$\begin{aligned} L_0 &= \sum_{n>0} n [N_{f_n} + N_{g_n} + N_{h_n} + N_{j_n}] + 1 \\ \lambda_0 &= \sum_{n>0} n [N_{u_n} + N_{v_n} + N_{w_n} + N_{z_n}] - 1 \\ K_0 &= \sum_{n>0} M [f_{-n} j_n + j_{-n} f_n] + (M f_0 + 2) j_0 \\ \kappa_0 &= \sum_{n>0} -n [z_{-n} u_n + u_{-n} z_n]. \end{aligned} \quad (16)$$

It follows immediately that all physical states must be products of matter and ghost ground states, and that they satisfy the matter zero-mode

projection

$$(Mf_0 + 2)j_0 |\text{phys}\rangle = 0. \quad (17)$$

Up to an additive constant ambiguity that arises from the addition of total derivatives linear in $\hat{g}^{++}$ or $\hat{X}^-$ to the action, f_0 and $-e^{\sqrt{2}M\tau}j_0$ are just the zero frequency components of the momenta conjugate to $\sqrt{2}\hat{X}^-/4\alpha'$ and $\hat{g}^{++}$, respectively. The condition (17) therefore has the same form as the projection onto the tachyon state in the conformal gauge, though here the generator of translations in $\hat{X}^+$ is tied through the gauge condition to the choice of world-sheet coordinates and so to the generator of translations in $\hat{g}^{++}$.

One possibility is that this is actually the correct spectrum of a consistent $1+1$ string theory, and that the special states are somehow artifacts of other methods of quantization. However, the fact that they are needed to obtain a consistent factorization of scattering amplitudes in [4], as well as their presence in the matrix models, makes this seem unlikely. The alternative is that there is some flaw or subtlety associated with this quantum theory which needs yet to be explained.

I again am indebted to J. Polchinski for useful discussions and for the use of his manuscript in preparation. I also thank R. Rudd for discussions. This research was supported in part by the Robert A. Welch Foundation, NSF Grant PHY 9009850, and the Texas Advanced Research Program.

References

- [1] P. Goddard and C. Thorn, Phys. Rev. Lett. B40 (1972) 235;
J.L. Gervais and B. Sakita, Nucl. Phys. B34 (1971) 477;
C.G. Callan, D. Friedan, E.J. Martinec and M.J. Perry, Nucl. Phys. B262 (1985) 593;
R.C. Myers, Phys. Lett. B199 (1987) 371
- [2] A.M. Polyakov, Mod. Phys. Lett. A2 (1987) 893;
V.G. Knizhnik, A.M. Polyakov, and A.B. Zamolodchikov, Mod. Phys. Lett. A3 (1988) 819
- [3] S. Mukherji, S. Mukhi and A. Sen, Phys. Lett. B266 (1991) 337;
U.H. Danielsson and D.J. Gross, Nucl. Phys. B366 (1991) 3;
A.M. Polyakov, Princeton preprint PUPT-1289, Lectures given at the Jerusalem winter school (1991);
I.R. Klebanov and A.M. Polyakov, Mod. Phys. Lett. A6 (1991) 3273
- [4] D. Minic and Z. Yang, Phys. Lett. B274 (1992) 27
- [5] E. Witten, Nucl. Phys. B373 (1992) 187;
D. Kutasov, E. Martinec and N. Seiberg, Phys. Lett. B276 (1992) 437
- [6] E. Smith, U.T. Austin preprint UTTG-36-91, to appear in Nucl. Phys. B
- [7] L.D. Faddeev and V.N. Popov, Phys. Lett. B25 (1967) 30
- [8] F. David, Mod. Phys. Lett. A3 (1988) 1651;
J. Distler and H. Kawai, Nucl. Phys. B321 (1989) 509;
J. Polchinski, Nucl. Phys. B324 (1989) 123
- [9] The author learned these methods from a manuscript on String Theory and Conformal Field Theory by J. Polchinski, in preparation.